

**Region 1,3 SITUATION STATUS (SIT/STAT) SUMMARY**  
**Bernalillo , Cibola, McKinley, Sandoval, San Juan and Valencia Counties**

Incident Name: H1N1 Region 1,3 Ready

Date & Time Prepared: 09/25/2009

SIT/STAT Type: Update

**SITUATION SUMMARY:**

Item	Status/Date	Update
Target Populations	Amended 9/25/09 See third bullet—revised to CDC written guidance	Initial delivery of vaccine expected to be 300,000 doses in NM. Initial priority populations are now: <ul style="list-style-type: none"> <li>• Pregnant women (unchanged)</li> <li>• Household contacts &amp; care givers of children &lt; 6 months (unchanged)</li> <li>• Health care workers and emergency medical personnel with direct patient contact</li> <li>• Children ages 6 – 59 months</li> <li>• Children ages 5 – 18 with underlying medical conditions that can result in complications from influenza</li> </ul>
Vaccine Clinical trials	Unchanged	Clinical trials are demonstrating that healthy adults will need only 1 dose. See vaccine administration section below
Vaccine Delivery	Updated 9/25/09 See number of doses	Delivery expected mid-October <i>initial shipment will be 20,000 to the state consisting of primarily FluMist—target population for this shipment will be healthy 2 – 4y os.</i> Providers, pharmacies and facilities encouraged to become direct ship sites information on DOH web page <a href="http://www.state.nm.us">www.state.nm.us</a> Vaccine will be available in the form of FluMist, thimerisol free, pre-filled syringes and multi-dose vials Initial delivery may not include vaccine for ages 6 mo-36 mo, even though included in priority group
Vaccine Administration	Unchanged	Dose information from vaccine inserts: <u>Children</u> Children 6 through 35 months of age should receive two 0.25 mL intramuscular doses approximately 1 month apart. (1) Children 36 months through 9 years of age should receive two 0.5 mL intramuscular doses approximately 1 month apart. (1) Children 10 years of age and older should receive a single 0.5 mL intramuscular dose. (1) The preferred sites for intramuscular injections are the anterolateral aspect of the thigh in infants or the deltoid muscle of the upper arm in toddlers and young children. The vaccine should not be injected into the gluteal region or into areas where there may be a major nerve trunk. <u>Adults</u> Persons 18 years of age and older should receive a single 0.5 mL intramuscular dose. In adults, the preferred site for intramuscular injection is the deltoid muscle.

<b>Prepared By:</b>	<b>Margy Wienbar</b>	<b>PH Region 1,3</b>		
		<b>505-841-4110</b>		

**Region 1,3 SITUATION STATUS (SIT/STAT) SUMMARY**  
**Bernalillo , Cibola, McKinley, Sandoval, San Juan and Valencia Counties**

		Confidential/Proprietary information The vaccine should not be injected into the gluteal region or into areas where there may be a major nerve trunk. See pdf file attached
County Stakeholder's Meetings	Completed	Bernalillo County 8/27/09 183 attendees Valencia County 9/9/09 39 attendees McKinley County 9/14/09 76 attendees Sandoval County 9/14/09 47 attendees San Juan County 9/16/09 95 attendees Cibola County n/a
Anti-virals	Updated 9/22/09	Updated CDC Guidance; Clarifies guidance for health care providers on early empiric use of antivirals. Early empiric treatment with oseltamivir or zanamivir should be considered for persons with suspected or confirmed influenza who are at higher risk for complications including: <ul style="list-style-type: none"> <li>• Children younger than 2 years old;</li> <li>• Persons aged 65 years or older</li> <li>• Pregnant women</li> <li>• Persons of any age with certain chronic medical or immunosuppressive conditions (see page 3); and,</li> <li>• Persons younger than 19 years of age who are receiving long-term aspirin therapy.</li> </ul> <a href="http://www.cdc.gov/h1n1flu/recommendations.htm">http://www.cdc.gov/h1n1flu/recommendations.htm</a>
Impact Summary	New 9/25/09	Nationally, deaths are lower and hospitalization rates are comparable to a typical flu season. NM is reporting "widespread" Influenza-like illness. To date there have been 5 deaths in NM due to H1N1. The most recent a 48 yo Bernalillo County man who had underlying health conditions. Severity of illness remains mild in most healthy individuals and most people are recovering without medical intervention.
CDC resources	Unchanged	Conference calls for local health departments initiated Web page <a href="http://www.flu.gov">www.flu.gov</a> consolidated resources for providers and the public Of note; Guidance for Small Businesses
Communications	Updated 9/25/09	Posters, bookmarks and immunization cards are in the process of being printed. These materials will be available for distribution through local public health. <i>Posters available on a limited basis being distributed through local public health offices or in Bernalillo County by calling Nicole Matthews at 841-4111</i>
Public Information Numbers	Updated 9/23/09	Nurse Advice Line available English and Spanish 1-877-725-2552—Reporting a 15 – 20% increase in volume Spanish information hotline 1-800-784-0394
Media	Unchanged	PSAs have been filmed and will be released

<b>Prepared By:</b>	<b>Margy Wienbar</b>	<b>PH Region 1,3</b>		
		<b>505-841-4110</b>		